

PYROL: Reg No. 7062 Act 36 of 1947

Contact insecticide containing Pyrethrin and Canola Oil

DIRECTIONS FOR USE: (use only as directed)

Broad spectrum

Shake well before use. Add sufficient water to spray tank and then add Pyrol, continue with agitation while adding additional water. Spray solution should be used within several hours of preparation.

Pyrol is suitable for indoor and outdoor use.

COMPATIBILITY:

Pyrol has been found to be compatible with most commonly used insecticides. Physical compatibility should first be checked by mixing the products in the recommended dosages in a small jar test. Then test the spray tank mix combinations for phytotoxicity on a sample of crops prior to use.

Do not mix with sulphur or copper compounds.

Do not apply in temperatures above 30 °C.

Ensure thorough wetting of the plant.

CROP	PESTS	DOSAGE	DIRECTIONS FOR APPLICATION
Ornamentals	Aphid	150 – 200 ml / 10 ℓ	Apply as a full cover spray. Ensure wetting of both top and underside of leaf. Repeat treatment as necessary. Established infestation will require frequent applications. Use higher dosage in times of higher infestation.
	Australian bug	150 – 200 ml / 10 ℓ	
Vegetables			
Cabbage	Turnip aphid	150 – 200 ml / 10 ℓ	Under controlled environments (eg tunnels, greenhouses etc) do a sampled, proportional jar test for phytotoxicity on crop, before spraying large areas. Recommend testing at the lowest recommended dosage for control.
Tomatoes	Green aphid	150 – 200 ml / 10 ℓ	
	Red spider mite	150 – 200 ml / 10 ℓ	
Citrus	Brown aphid	150 – 200 ml / 10 ℓ	

PYROL: Reg No. 7062 Act 36 of 1947

Contact insecticide containing Pyrethrin and Canola Oil

GEBRUIKSAANWYSINGS: (Gebruik slegs soos aangedui.)

Broad spectrum

Skud goed voor gebruik.

Voeg voldoende water in spuitenk en voeg dan Pyrol by. Hou aan met meng terwyl addisionele water bygegooi word. Die spuitoplossing moet binne 'n paar uur na voorbereiding gebruik word. Pyrol is gepas vir binne- of buitemuurse gebruik.

VERENIGBAARHEID:

Daar is bevind dat Pyrol met die meeste algemeen-gebruikte insekdoders gemeng kan word. Fisiese mengbaarheid behoort vooraf getoets te word deur die produkte in die aanbevole dosisse in 'n klein houer te toets.

Om fitotoksisiteit te bepaal, behoort spuitenk meng-kombinasies op 'n monster van gewasse getoets te word.

Moenie met swael of koper verbindings gemeng word nie.

Moenie in temperature bo 30 °C toedien nie.

Maak seker dat die hele plant deeglik benat word.

Gewas	Plaag	Dosis	Aanwysings vir toediening
Sierplante	Plantluis	150 – 200 ml / 10 ℓ	Tydens toediening moet verseker word dat die plante ten volle deur die spuitstof gedek word. Verseker dat blare aan die bokant en onderkant benat word. Herhaal behandeling soos benodig. Gevestigde infestaties sal verskeie aanwendings benodig. Gebruik hoër dosisse gedurende hoë infestaties.
	Australiese luis	150 – 200 ml / 10 ℓ	
Groente			
Kool	Plantluis	150 – 200 ml / 10 ℓ	Onder gekontroleerde omgewings (tonnels, kweekhuise ens) behoort kwekers 'n monster van die gewas te toets vir plant/fitotoksisiteit, voor groot areas bespuit word. Toets gewasse teen die laagste aanbevole dosis.
Tamaties	Plantluis	150 – 200 ml / 10 ℓ	
	Rooispinmyt	150 – 200 ml / 10 ℓ	
Sitrus	Plantluis	150 – 200 ml / 10 ℓ	